



The following information is provided for the 51-60 range of the DWR
 H (CWR, LM, VCR, ECGL, ETLR, 1979. USFWS, SRF,
 F S/OBS-79/31, 131). The following information is provided for the NRI
 DWR range.




R 51. K :Pl r V :N , r O r C 50
 The r S J M . (G C , Cl -
 r ; Pl R.M. H r)



R 52. K y :Pl r V :N r O l r C 50
 Tl l w r r l ; l y
 r l r r r (*Kochia scopari*), l
 (*Rumex maritimu*), (*Chenopodium glaucu*). (S
 C y, N r l D ; A 1961; Pl y R.E. S w r)




R 53. K :Pl r V :N , r O l r C 50
A r r l (*Alisma plantago-aquatic*) r l l
L l l . (Bl C y, N r l D ; J l y 1970; Pl l J.T.




R 54. K y :Pl r V :N , r O l r C 50
D F y : l w r F w (*Ranunculus trichophyllu*). (S
C y, N r l D ; A 1966; Pl y R. E. S w r)




R 55. K : Pl r V : N , r O l r C 50
 T- l (Sphagnum). S r l
 r r (Cladonia), l l (Chamaedaphne calyculat), F w-
 rr y (Empetrum nigrum), F r (Eriophorum). (C l
 Il I r l Pr , M C ; J 1976; Pl y V. C r r)



R 56. K :Pl r V :N , r O l r C 50
P (S l .) l .S r l
(Carex rariflora, C. aquatili), F r (Eriophorum russeoli),
r r (Cladina). l r r , l r
F r l l 30%. M F r 100% l r . (N r l A R -
r r , K w D l , A ; J l y 1985; Pl l F.C. G l)




R 57 K :Pl r V :E r C 51
D F y :D F y C l (*Typha latifolia*). Pr
r r w .N l l F-F r l
r l y. (K C y, M ; A r l 1978; Pl y P. B. R)



R 58 K y Pl r V : E C 51
D B : B l r l (*Scirpus robustu*) H r l r l (*Scirpus*
acutu). (S C y, N r l D ; A 1962; Pl- y R.E. S w r)



Figure D-59. Key Plant Species: *Cladium jamaicens* (Water Hyacinth) in a Wetland Habitat. (Data from U.S. Fish and Wildlife Service, 1975; Plant Species List).



R 60 K y :Pl r V :E r C 51
D F y :S (*Carex lasiocarp*). S r l
(*Carex lacustris*, *C. rostrat*), w r r w (*Polygonum amphibiu*), l -
r w r (*Utricu-laria macrorhiz*), l (*Calamagrostis canadensi*),
w (*Potamogeton gramineu*). (Cl- w N l Fr , Bl r
C y, M ; J 1972; Pl- y J.H. R)